

In Vitro Anti-Inflammatory Potential of Buton Forest Honey through Nitric Oxide Inhibition in RAW 264.7 Macrophage Cells

Hariana¹, Wahyuni², Suryani², Ari Sartinah², Maulidyah³, Agung Wibawa Mahatva Yodha⁴, La Ode Muhammad Julian Purnama⁵, Adryan Fristiohady^{2*}

¹ Program Studi Magister Farmasi, Fakultas Farmasi, Universitas Halu Oleo, 93232, Sulawesi Tenggara, Indonesia

² Jurusan Farmasi, Fakultas Farmasi, Universitas Halu Oleo, Kampus Hijau Bumi Tridharma, Anduonohu, Kec. Kambu, Kota Kendari, Sulawesi Tenggara 93232, Indonesia

³ Jurusan Kimia, Fakultas Matematika dan Ilmu Pengetahuan Alam, Universitas Halu Oleo, Kampus Hijau Bumi Tridharma, Anduonohu, Kec. Kambu, Kota Kendari, Sulawesi Tenggara 93232, Indonesia

⁴ Program Studi Diploma III Farmasi, Politeknik Bina Husada, Kota Kendari, Sulawesi Tenggara 93232, Indonesia

⁵ Faculty of Pharmacy, Thammasat University Research Unit in Drug, Health Product Development and Application (DPH-DA), Thammasat University, Khlong Luang, 12120, Pathum Thani, Thailand.

* Corresponding Author. E-mail: adryanfristiohady@uho.ac.id

Received: 17 October 2025 / Accepted: 28 March 2026 / Published: 30 March 2026

ABSTRACT: Inflammation is a physiological defense response to injury, infection, or stress, mediated by various signaling molecules, including nitric oxide (NO). However, excessive NO production can contribute to tissue damage and the progression of inflammatory conditions. Honey, a natural product derived from floral nectar by bees, contains diverse bioactive compounds with potential pharmacological properties, including anti-inflammatory activity. Buton Regency, located in Southeast Sulawesi, is a major producer of forest honey, making it a promising source of bioactive honey with therapeutic potential. This study aimed to evaluate the in vitro anti-inflammatory activity of Buton forest honey by assessing its ability to inhibit NO production in lipopolysaccharide (LPS)-stimulated RAW 264.7 macrophage cells. The results demonstrated that Buton forest honey significantly suppressed NO production in a concentration-dependent manner. At concentrations of 500, 250, 125, 62.5, 31.25, 15.63, 7.81, and 3.91 mg/L, the percentages of inhibition were $94.69 \pm 2.94\%$, $74.03 \pm 2.43\%$, $67.40 \pm 2.34\%$, $62.72 \pm 3.09\%$, $51.80 \pm 4.73\%$, $44.40 \pm 4.73\%$, $19.06 \pm 1.79\%$, and $12.82 \pm 0.68\%$, respectively. Statistical analysis using one-way ANOVA followed by post hoc testing revealed significant differences among the tested concentrations ($p < 0.05$). In conclusion, Buton forest honey exhibits significant anti-inflammatory activity by inhibiting NO production in macrophage cells, suggesting its potential as a natural source for the development of anti-inflammatory agents.

KEYWORDS: Anti-inflammatory; honey; macrophage cell; nitric oxide; protective.

1. INTRODUCTION

Inflammation is a fundamental physiological defense mechanism that occurs in response to infection, injury, or physiological stress. This process involves a complex interplay of immune cells and chemical mediators that function to restore tissue homeostasis. However, when inflammation becomes excessive or chronic, it can lead to tissue damage and contribute to the pathogenesis of various chronic diseases, including arthritis, diabetes mellitus, atherosclerosis, and cancer (Fristiohady et al., 2019; Fristiohady A, 2020; Yodha et al., 2024; Fawwaz et al., 2024). Among the key mediators involved in inflammation is nitric oxide (NO), a signaling molecule produced by inducible nitric oxide synthase (iNOS) in activated macrophages. While NO plays an important role in host defense, its overproduction has been associated with the amplification of inflammatory responses and cellular damage. Therefore, the inhibition of NO synthesis has emerged as an important therapeutic target in the development of anti-inflammatory agents (Poulsen-Silva et al., 2023; Sukmawati et al., 2023).

In recent years, increasing interest in natural-based therapies has highlighted honey as a promising candidate for anti-inflammatory applications. In addition to its nutritional value, honey contains a wide range of bioactive compounds, including flavonoids, phenolic compounds, terpenoids, and organic acids, which contribute to its diverse biological activities such as antioxidant, antibacterial, and anti-inflammatory effects (Alqarni et al., 2014; Hariana & Fristiohady, 2025; Hulea et al., 2022). Among the various types of honey, forest honey possesses distinctive characteristics due to its production by wild bees that forage on a diverse array of forest flora. This results in a more complex chemical composition and potentially enhanced bioactivity. Furthermore, variations in nectar sources and environmental conditions contribute to the unique pharmacological properties of honey from different geographical regions (Dewantara, 2021; Lawag et al., 2023; Ndungu et al., 2024).



The investigation of anti-inflammatory mechanisms requires appropriate experimental models, and the RAW 264.7 macrophage cell line is widely used as a standard *in vitro* system for studying inflammatory responses. Upon stimulation with lipopolysaccharide (LPS), these cells produce various inflammatory mediators, including nitric oxide (NO), tumor necrosis factor- α (TNF- α), and interleukin-6 (IL-6) (Faris, 2019; Pham *et al.*, 2021). Consequently, the inhibition of NO production in LPS-stimulated RAW 264.7 cells is commonly employed as a primary indicator of *in vitro* anti-inflammatory activity. Previous studies have demonstrated that honey can suppress the production of these mediators by modulating inflammatory signaling pathways in macrophages (Yuandani *et al.*, 2016).

Southeast Sulawesi Province is one of the regions in Indonesia with significant potential for forest honey production (BPS, 2024; Nurhasanah Sari *et al.*, 2018). In particular, Buton Regency is characterized by well-preserved tropical forests and a high diversity of nectar-producing plants, which contribute to the unique chemical profile and bioactive composition of its forest honey (Uji *et al.*, 2007). Although several studies have reported the anti-inflammatory activity of honey from various regions through the inhibition of inflammatory mediators such as nitric oxide (Romário-Silva *et al.*, 2022), research specifically investigating the anti-inflammatory potential of forest honey from Buton Regency remains limited.

Therefore, this study aims to evaluate the *in vitro* anti-inflammatory activity of Buton forest honey by examining its ability to inhibit nitric oxide production in LPS-stimulated RAW 264.7 macrophage cells. This study is expected to provide new scientific insights into the pharmacological potential of forest honey from Southeast Sulawesi as a natural anti-inflammatory agent.

2. EXPERIMENTAL SECTION

2.1. Sample Preparation

Forest honey samples were collected from natural forest areas in Buton Regency, Southeast Sulawesi, Indonesia (geographical coordinates: 5°17'9.53"S, 122°46'38.39"E; -5.285981, 122.777331). The honey was identified as forest honey based on its physicochemical characteristics and confirmation by local apiarists. A 1-gram sample of this forest honey was subjected to sonication using a sonicator (Elmazonik®, India) with 10 mL of 70% ethanol (Onemed®, Indonesia) for 30 minutes at room temperature to ensure thorough mixing. Following homogenization, the mixture was filtered through filter paper, transferred into a 100 mL volumetric flask, and diluted with ethanol up to the calibration mark, resulting in a final concentration of 10,000 mg/L. The prepared sample was then stored at 2–4°C until further analysis for its anti-inflammatory activity (Hulea *et al.*, 2022).

2.2. Anti-inflammatory Activity Assay Using RAW 264.7 Macrophage Cells

RAW 264.7 cells were seeded in 96-well plates at a density of 2×10^5 cells/well (100 μ L) and incubated for 24 hours at 37 °C in a humidified atmosphere containing 5% CO₂. After incubation, the culture medium was removed, and the cells were washed twice with PBS. The cells were then treated with DMEM medium containing lipopolysaccharide (LPS, 1 μ g/mL) to induce nitric oxide (NO) production, followed by the addition of forest honey samples at various concentrations ranging from 3.91 to 500 μ g/mL. After incubation for 18–24 hours, 50 μ L of the supernatant was collected and mixed with 50 μ L of Griess reagent, consisting of 1% sulfanilamide in 2.5% phosphoric acid and 0.1% naphthylethylenediamine. The mixture was incubated for 10 minutes at room temperature, and the absorbance was measured at 540 nm using a microplate reader.

The nitrite concentration was determined based on a standard NaNO₂ calibration curve, while the percentage of NO inhibition was calculated using the following formula:

$$\text{Inhibition (Anti – inflammatory Activity)NO} = \frac{\text{Absorbance of control} - \text{Absorbance of Sample}}{\text{Absorbance of control}} \times 100 \text{ [1]}$$

The control absorbance was obtained from LPS-treated cells minus the absorbance of untreated cells, whereas the sample absorbance was calculated from LPS + sample-treated cells minus the absorbance of untreated cells (Ruangnoo *et al.*, 2012).

2.5. Data Analysis

Statistical analysis was performed using SPSS software through a one-way analysis of variance (ANOVA), followed by Tukey's post hoc test to determine significant differences among groups. A p-value of < 0.05 was considered the threshold for statistical significance.

3. RESULTS AND DISCUSSION

The anti-inflammatory activity of forest honey from Buton was evaluated by assessing its ability to inhibit nitric oxide (NO) production in lipopolysaccharide (LPS)-stimulated RAW 264.7 macrophage cells. Macrophages were selected as the experimental model due to their central role in inflammatory responses, as they produce a variety of mediators, including NO, prostaglandin E₂ (PGE₂), interleukin-6 (IL-6), and tumor necrosis factor- α (TNF- α), upon activation. Stimulation with LPS activates Toll-like receptor 4 (TLR4), which subsequently triggers intracellular signaling

pathways such as nuclear factor-kappa B (NF-κB) and mitogen-activated protein kinase (MAPK), leading to increased expression of the inducible nitric oxide synthase (iNOS) gene (Faris, 2019; Sukmawati *et al.*, 2023).

The iNOS enzyme catalyzes the conversion of L-arginine into NO, which is further oxidized into more stable metabolites, namely nitrite (NO₂⁻) and nitrate (NO₃⁻). In this study, NO production was quantified indirectly by measuring nitrite levels using the Griess assay. Prednisolone, a synthetic corticosteroid with well-established anti-inflammatory properties, was used as a positive control. This compound exerts its effects by suppressing the expression of pro-inflammatory enzymes such as iNOS and cyclooxygenase-2 (COX-2), as well as reducing the production of pro-inflammatory cytokines, including TNF-α, IL-6, and PGE₂, thereby serving as a reference standard for evaluating the efficacy of Buton forest honey (Hussein *et al.*, 2012). The results of NO inhibition, expressed as percentage inhibition of cell-mediated NO production, are presented in **Table 1** and **Figure 1**.

Table 1. Percentage of nitric oxide (NO) inhibition by Buton forest honey and prednisolone in LPS-stimulated RAW 264.7 macrophage cells.

Concentrations (mg/L)	Inhibition (%)	
	Buton forest honey	Prednisolone
500	94.69 ± 2.94	100.03 ± 1.79
250	74.03 ± 2.43	97.42 ± 3.38
125	67.40 ± 2.34	94.30 ± 6.51
62.5	62.72 ± 3.09	79.09 ± 2.34
31.25	51.80 ± 4.73	73.25 ± 2.34
15.625	44.40 ± 4.73	66.62 ± 2.94
7.8125	19.05 ± 1.79	44.79 ± 3.76
3.90625	12.82 ± 0.68	29.97 ± 0.68




Figure 1. Percentage of NO production inhibition by forest honey from Konawe in RAW 264.7 macrophage cells.

The graph illustrates the percentage of nitric oxide (NO) inhibition in LPS-stimulated RAW 264.7 macrophage cells following treatment with Buton forest honey, compared with prednisolone as a positive control, across a concentration range of 0–500 μg/mL. The horizontal axis represents the sample concentration (μg/mL), while the vertical axis indicates the percentage of NO inhibition. Prednisolone exhibited a rapid and pronounced inhibitory effect, achieving approximately 50–70% inhibition at lower concentrations and approaching near-complete inhibition at higher concentrations (250–500 μg/mL). In contrast, Buton forest honey demonstrated a clear concentration-dependent inhibitory effect, with NO inhibition gradually increasing as the concentration increased. Maximum inhibition, approaching 90–100%, was observed only at the highest concentration (500 μg/mL). The comparatively slower response

of Buton forest honey suggests a milder or more gradual mechanism of action, likely associated with the cumulative effects of its bioactive constituents.

Statistical analysis was performed using SPSS software, employing one-way ANOVA followed by Tukey's post hoc test to assess differences among treatment groups. The results revealed statistically significant differences ($p < 0.05$) between the control group and honey-treated groups at concentrations of 250 and 500 $\mu\text{g/mL}$, indicating effective suppression of NO production in a dose-dependent manner. Prednisolone exhibited significantly greater inhibitory activity ($p < 0.01$) compared to all honey concentrations, consistent with its role as a synthetic corticosteroid with potent and rapid anti-inflammatory effects.

Despite its lower potency relative to prednisolone, Buton forest honey demonstrated a significant and measurable inhibitory effect on NO production, underscoring its potential as a natural anti-inflammatory agent. The observed concentration-dependent inhibition aligns with pharmacological principles, where increased concentrations of active compounds result in enhanced biological effects. This activity is likely attributable to bioactive compounds such as flavonoids and phenolic constituents, which have been reported to suppress inflammatory signaling pathways, including nuclear factor-kappa B (NF- κB) activation and inducible nitric oxide synthase (iNOS) expression in RAW 264.7 macrophages (Faris, 2019; Pham et al., 2021).

In conclusion, although prednisolone demonstrated a faster and more potent inhibitory effect, Buton forest honey exhibited significant, dose-dependent anti-inflammatory activity. These findings support the hypothesis that its anti-inflammatory effects are biologically relevant and concentration-dependent, reinforcing its potential as a natural alternative or complementary agent to synthetic anti-inflammatory drugs.

This study also holds important scientific and practical implications. Excessive nitric oxide production is a key contributor to chronic inflammatory conditions (Fristiody et al., 2025; Hariana et al., 2025), and the significant inhibition observed in this study highlights the therapeutic potential of Buton forest honey. Furthermore, these findings support the valorization of Indonesian local biological resources not only as economic commodities but also as promising sources of nutraceuticals and complementary therapies. Future studies are warranted to further elucidate the molecular mechanisms underlying the anti-inflammatory effects of forest honey, particularly those involving nitric oxide regulation and iNOS-mediated pathways.

4. CONCLUSION

The findings of this study demonstrate that forest honey from Buton exhibits significant anti-inflammatory activity through the inhibition of nitric oxide (NO) production in LPS-stimulated RAW 264.7 macrophage cells. This inhibitory effect was concentration-dependent, with higher concentrations resulting in greater suppression of NO production. Statistical analysis using one-way ANOVA followed by Tukey's post hoc test confirmed a significant reduction in NO levels at concentrations of 250 and 500 $\mu\text{g/mL}$ ($p < 0.05$), indicating the biological relevance of the observed activity. Although prednisolone, as a synthetic corticosteroid, produced a more rapid and potent inhibitory effect, Buton forest honey demonstrated a substantial and measurable anti-inflammatory response. This activity may be attributed to the presence of bioactive compounds such as flavonoids and phenolic constituents, which are known to modulate key inflammatory pathways, including nuclear factor-kappa B (NF- κB) signaling and inducible nitric oxide synthase (iNOS) expression. Overall, these findings suggest that Buton forest honey holds promise as a natural anti-inflammatory agent and may serve as a complementary alternative to conventional synthetic drugs.

Acknowledgments: The authors gratefully acknowledge the Directorate of Research and Community Service, Directorate General of Higher Education, Research, and Technology, Ministry of Education, Research, and Technology of the Republic of Indonesia for supporting this research through the Postgraduate Thesis Research Grant Scheme (PPS-PTM) 2025 (Master Contract No. 068/C3/DT.05.00/PL/2025 and Sub-Contract No. 07/UN29.20/PG/2025).

Author contributions: All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Conflict of interest: The authors declared no conflict of interest.

Ethical Approval: Not applicable

REFERENCES

- Alqarni, A. S., Owayss, A. A., Mahmoud, A. A., & Hannan, M. A. Mineral content and physical properties of local and imported honeys in Saudi Arabia. *Journal of Saudi Chemical Society*, 2014; 18(5), 618–625. <https://doi.org/10.1016/j.jscs.2012.11.009>
- BPS. (2024). Statistik Produksi Kehutanan.
- Dewantara, I. Pengelolaan Madu Lalau Oleh Masyarakat Desa Nanga Lauk Kecamatan Embaloh Hilir Kabupaten Kapuas Hulu Forest Honey Management by the Community in Nanga Lauk Village Embaloh District of Kapuas Hulu Regency, 2021; 9(1).
- Fawwaz, M., Purwono, B., Sidiq, Y., Arwansyah, Arsul, M. I., Fitriana & Baits, M. Anti-inflammatory, Antioxidant, and Antibacterial Activities with Molecular Docking Studies of Vitex Trifolia L. Targeting Human COX-2 and Peroxiredoxin-5. *ChemistrySelect*, 2024; 9(39), e202403834.

- Faris, M. Potensi Immunodulator Ekstrak Cengkeh Pada Kadar Limfosit Dan Makrofag Sebagai Mekanisme Pertahanan Tubuh. 2019.
- Fristiohady, A, Wahyuni, *et al.* In Vitro Anti-inflammatory Activity of *Elingera elatior* (Jack) R.M. Smith by Hrbc Membrane Stabilization Method. *Jurnal Ilmu Kefarmasian Indonesia*, 2020.
- Fristiohady, A., Ode Muhammad Fitrawan, L., Subhiaty Idrus, L., & Wibawa Mahatva Yodha, A. Potensi Radical Scavenging Dan Aktivitas Anti-Inflamasi Melalui Penghambatan Nitric Oxide Dari Madu Hutan Asal Kabupaten Konawe. *J. Sains Dan Teknologi Pangan*, 2025; 10(4), 8670–8884.
- Fristiohady, A., Wahyuni, Malik, F., Purnama, L. O. M. J., Sadarun, B., & Sahidin. Anti-inflammatory activity of marine sponge callyspongia sp. And its acute toxicity. *Asian Journal of Pharmaceutical and Clinical Research*, 2019; 97–100. <https://doi.org/10.22159/ajpcr.2019.v12i12.34737>
- Hariana, & Fristiohady, A. Tinjauan Farmakologi Madu: In Vitro dan In Vivo. Yogyakarta. 2025.
- Hariana, H., Wahyuni, W., Irnawati, I., Ode, L., Fitrawan, M., Munasari, D., Wibawa, A., Yodha, M., Purnama, M. J., & Fristiohady, A. Metabolite Profiling and Anti-Inflammatory Potential of East Kolaka Forest Honey: Targeting Protein Denaturation through GC-MS/MS Characterization. *Jurnal Mandala Pharmacoon Indonesia (JMPI)*, 2025; 11(2), 362–373. <https://doi.org/10.35311/jmpi.v11i2.860>
- Hulea, A., Obiștioiu, D., Cocan, I., Alexa, E., Negrea, M., Neacșu, A. G., Hulea, C., Pascu, C., Costinar, L., Iancu, I., Tîrziu, E., & Herman, V. Diversity of Monofloral Honey Based on the Antimicrobial and Antioxidant Potential. *Antibiotics*, 2022; 11(5). <https://doi.org/10.3390/antibiotics11050595>
- Hussein, S. Z., Mohd Yusoff, K., Makpol, S., & Mohd Yusof, Y. A. Gelam honey inhibits the production of proinflammatory mediators NO, PGE 2, TNF- α , and IL-6 in carrageenan-induced acute paw edema in rats. *Evidence-Based Complementary and Alternative Medicine*, 2012. <https://doi.org/10.1155/2012/109636>
- Lawag, I. L., Islam, M. K., Sostaric, T., Lim, L. Y., Hammer, K., & Locher, C. Antioxidant Activity and Phenolic Compound Identification and Quantification in Western Australian Honeys. *Antioxidants*, 2023; 12(1). <https://doi.org/10.3390/antiox12010189>
- Ndungu, N. N., Kegode, T. M., Kurgat, J. K., Baleba, S. B. S., Cheseto, X., Turner, S., Tasse Taboue, G. C., Kasina, J. M., Subramanian, S., & Nganso, B. T. Bio-functional properties and phytochemical composition of selected *Apis mellifera* honey from Africa. *Heliyon*, 2024; 10(10). <https://doi.org/10.1016/j.heliyon.2024.e30839>
- Nurhasanah Sari, D., Rifkah Ansyarif, A., Syekh Yusuf Al Makassar Gowa, U., Melati No, J., & Bonto-Bontoa Kec Somba Opu Kab Gowa, K. Karakteristik Madu Hutan Lebah Apis Dorsata Daerah Sulawesi Tenggara Ditinjau Dari Sifat Fisika-Kimia. In *Cokroaminoto Journal of Chemical Science*, 2018; 5(2).
- Pham, T. M., Erichsen, J. L., Kowal, J. M., Overgaard, S., & Schmal, H. Elevation of pro-inflammatory cytokine levels following intra-articular fractures a systematic review. *Cells*, 2021; 10(4). <https://doi.org/10.3390/cells10040902>
- Poulsen-Silva, E., Gordillo-Fuenzalida, F., Velásquez, P., Llancahuén, F. M., Carvajal, R., Cabaña-Brunod, M., & Otero, M. C. Antimicrobial, Antioxidant, and Anti-Inflammatory Properties of Monofloral Honeys from Chile. In *Antioxidants*, 2023; 12(9). Multidisciplinary Digital Publishing Institute (MDPI). <https://doi.org/10.3390/antiox12091785>
- Romário-Silva, D., Lazarini, J. G., Franchin, M., de Alencar, S. M., & Rosalen, P. L. Brazilian Organic Honey from Atlantic Rainforest Decreases Inflammatory Process in Mice. *Veterinary Sciences*, 2022; 9(6). <https://doi.org/10.3390/vetsci9060268>
- Ruangnoo, S., Jaiaree, N., Makchuchit, S., Panthong, S., Thongdeeying, P., & Itharat, A. *Original article An in vitro inhibitory effect on RAW 264* (Number 7). 2012.
- Sukmawati, Musfiroh, I., Muchtaridi, Fristiohady, A., & Ikram, N. K. K. Anti-Inflammatory Activity of Quts Al-Hindi Extract (*Saussurea lappa*) Using Erythrocyte Membrane Stability and Prediction of Its Selectivity for COX-2 and INOS Enzymes. *Journal of Hunan University Natural Sciences*, 2023; 50(10). <https://doi.org/10.55463/issn.1674-2974.50.10.3>
- Uji, T., Peneliti, F. I. W., Penelitian, P., Lembaga, B., & Pengetahuan Indonesia, I. Keanekaragaman Jenis Tumbuhan Di Cagar Alam Kakenauwe Dan Suaka Margasatwa Lambusango, Pulau Buton Sulawesi Tenggara. 2007.
- Yodha, A. W. M., Badia, E., Musdalipah, Reymon, Fauziah, Y., Fusvita, A., Arfan, Wahyuni, & Sahidin. Secondary Metabolite Compounds from *Alpinia monopleura* Extract and Evaluation of Anti-Inflammatory Activity based on In Vitro and In Silico Studies. *Hayati Journal of Biosciences*, 2024; 31(6), 1154–1164. <https://doi.org/10.4308/hjb.31.6.1154-1164>
- Yuandani, Jantan, I., Ilangkovan, M., Husain, K., & Chan, K. M. Inhibitory effects of compounds from *Phyllanthus amarus* on nitric oxide production, lymphocyte proliferation, and cytokine release from phagocytes. *Drug Design, Development and Therapy*, 2016; 10, 1935–1945. <https://doi.org/10.2147/DDDT.S105651>

This is an open access article which is publicly available on our journal's website under Institutional Repository at <https://www.umi.ac.id/id/>